

Native Human Apolipoprotein A2 HDL

Catalog No.	CSI19782A CSI19782B	Quantity:	1.0 mg 2.5 mg
Description:	25 percent of Apo A in HDL is Apo AII. Levels of Apo AII in normal plasma range from 30 -50 mg per 100 ml. The physiological role of the protein is unknown. Alcohol consumption, however, increases Apo AII levels. Prepared from plasma shown to be non reactive for HbsAG, anti-HCV, anti-HBc, and negative for anti-HIV 1 & 2 by FDA approved tests. >95% pure by SDS-PAGE		
Concentration:	10.0 mg/ml		
Source:	Human plasma		
Molecular Weight:	17.4 kDa		
Formulation:	Liquid in 0.01M Ammonium Bicarbonate, pH 7.4.		
Purity:	>95% pure by SDS-PAGE.		
Storage & Stability:	When stored at -80°C, product is stable for 3 years from date of delivery. Keep at a concentration ≥ 1 mg/ml. Avoid repeated freeze-thaw cycles.		
Certification:	Prepared from plasma shown to be non reactive for HbsAG, anti-HCV, anti-HBc, and negative for anti-HIV 1 & 2 by FDA approved tests. >95 % pure by SDS-PAGE.		

NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.

