

Synonym


Spike,S protein RBD,Spike glycoprotein Receptor-binding domain,S glycoprotein RBD,Spike protein RBD,COVID-19

Source

Biotinylated SARS-CoV-2 S protein RBD, His,Avitag (SPD-C82E9) is expressed from human 293 cells (HEK293). It contains AA Arg 319 - Phe 541 (Accession # [QHD43416.1](#)).

Predicted N-terminus: Arg 319

Molecular Characterization



This protein carries a polyhistidine tag at the C-terminus, followed by an Avi tag.

The protein has a calculated MW of 28.2 kDa. The protein migrates as 33-36 kDa under reducing (R) condition (SDS-PAGE) due to glycosylation.

Biotinylation

Biotinylation of this product is performed using Avitag™ technology. Briefly, the single lysine residue in the Avitag is enzymatically labeled with biotin.

Biotin:Protein Ratio

The biotin to protein ratio is *0.5-1* as determined by the HABA assay.

Endotoxin

Less than 1.0 EU per µg by the LAL method.

Purity

>95% as determined by SDS-PAGE.

>90% as determined by SEC-MALS.

Formulation

Lyophilized from 0.22 µm filtered solution in PBS, pH7.4. Normally trehalose is added as protectant before lyophilization.

Contact us for customized product form or formulation.

Reconstitution

Please see Certificate of Analysis for specific instructions.

For best performance, we strongly recommend you to follow the reconstitution protocol provided in the CoA.

Storage


For long term storage, the product should be stored at lyophilized state at -20°C or lower.

Please avoid repeated freeze-thaw cycles.

This product is stable after storage at:

- -20°C to -70°C for 12 months in lyophilized state;
- -70 °C for 12 months under sterile conditions after reconstitution.


SDS-PAGE



Biotinylated SARS-CoV-2 S protein RBD, His,Avitag on SDS-PAGE under reducing (R) condition. The gel was stained overnight with Coomassie Blue. The purity of the protein is greater than 95%.


Bioactivity-ELISA

SEC-MALS




The purity of Biotinylated SARS-CoV-2 S protein RBD, His,Avitag (Cat. No. SPD-C82E9) was more than 90% and the molecular weight of this protein is around 33-43 kDa verified by SEC-MALS.

[Report](#)




Immobilized Human ACE2, Fc Tag (Cat. No. [AC2-H5257](#)) at 2 µg/mL (100 µL/well) can bind Biotinylated SARS-CoV-2 S protein RBD, His,Avitag (Cat. No. [SPD-C82E9](#)) with a linear range of 2-78 ng/mL (QC tested).



Immobilized Biotinylated SARS-CoV-2 S protein RBD, His,Avitag (Cat. No. [SPD-C82E9](#)) at 1 µg/mL (100 µL/well) on streptavidin precoated (0.5 µg/well) plate, can bind Anti-SARS-CoV-2 Neutralizing Antibody, Human IgG1 (Cat. No. [SAD-S35](#)) with a linear range of 0.1-3 ng/mL (Routinely tested).

Bioactivity-BLI



Loaded Biotinylated SARS-CoV-2 S protein RBD, His,Avitag (Cat. No. SPD-C82E9) on SA Biosensor, can bind Human ACE2, His Tag (Cat. No. AC2-H52H8) with an affinity constant of 1.14 nM as determined in BLI assay (ForteBio Octet Red96e) (Routinely tested).

Background

It's been reported that SARS-CoV-2 can infect the human respiratory epithelial cells through interaction with the human ACE2 receptor. The spike protein is a large type I transmembrane protein containing two subunits, S1 and S2. S1 mainly contains a receptor binding domain (RBD), which is responsible for recognizing the cell surface receptor. S2 contains basic elements needed for the membrane fusion. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

References

- (1) [Wan Y, et al. J Virol. 2020. pii: JVI.00127-20.](#)
- (2) [Benvenuto D, et al. J Med Virol. 2020.](#)
- (3) [Chang CY, et al. AMB Express. 2020. 10\(1\):20.](#)

Please contact us via TechSupport@acrobiosystems.com if you have any question on this product.