

MAB040Hu23

**Monoclonal Antibody to Vimentin (VIM)** 

Organism Species: Homo sapiens (Human)

Instruction manual

FOR RESEARCH USE ONLY
NOT FOR USE IN CLINICAL DIAGNOSTIC PROCEDURES

12th Edition (Revised in Aug, 2016)



## [PROPERTIES]

**Source:** Monoclonal antibody preparation

Host: Mouse

Antibody isotype: IgG2b Kappa

**Purification:** Protein A + Protein G affinity chromatography

Clone number: C5

Traits: Liquid

Concentration: 1mg/ml

**UOM:** 200µl

Cross Reactivity: Rabbit; Canine; Porcine; Bovine; Caprine; Ovine; Gallus

Applications: WB; IHC; ICC; IP.

### [ IMMUNOGEN ]

Immunogen: Recombinant VIM (Ser2~Glu466) expressed in E.coli

Accession No.: RPB040Hu01

#### [ APPLICATIONS ]

Western blotting: 0.5-2µg/mL

Immunohistochemistry: 5-20µg/mL

Immunocytochemistry: 5-20µg/mL

Optimal working dilutions must be determined by end user.

#### [FORMULATION]

Form & Buffer: Supplied as solution form in 0.01M PBS, pH7.4, containing 0.05% Proclin-300,

50% glycerol.

#### [ STORAGE AND STABILITY ]

Storage: Avoid repeated freeze/thaw cycles.

Store at 4°C for frequent use.


Aliquot and store at -20°C for 12 months.

Stability Test: The thermal stability is described by the loss rate. The loss rate was determined


# Coud-Clone Corp.

by accelerated thermal degradation test, that is, incubate the protein at 37°C for 48h, and no obvious degradation and precipitation were observed. The loss rate is less than 5% within the expiration date under appropriate storage condition.

## [ IDENTIFICATION ]



Western Blot; Sample: Lane1: Canine
Heart lysate; Lane2: Gallus Heart
lysate; Lane3: Porcine Heart lysate;
Lane4: Bovine Heart lysate; Lane5:
Caprine Heart lysate; Lane6: Caprine
Heart lysate.



DAB staining on IHC-P; Sample:
Human Breast cancer Tissue; Primary
Ab: 10µg/ml Mouse Anti-Human VIM
Antibody Second Ab: 2µg/mL HRPLinked Caprine Anti-Mouse IgG
Polyclonal Antibody (Catalog:
SAA544Mu19)

# [ IMPORTANT NOTE ]

The kit is designed for research use only, we will not be responsible for any issue if the kit was used in clinical diagnostic or any other procedures.