

**CD57 Antibody**  
**Mouse Monoclonal Antibody (Mab)**  
**Catalog # AD80190**

**Specification**

**CD57 Antibody - Product info**

Application	<b>IHC</b>
Primary Accession	<a href="#">Q9P2W7</a>
Reactivity	<b>Human</b>
Host	<b>Mouse</b>
Clonality	<b>Monoclonal</b>
Isotype	<b>IgM</b>
Calculated MW	<b>38256</b>

**CD57 Antibody - Additional info**

Gene ID	<b>27087</b>
Gene Name	<b>B3GAT1</b> ( <a href="#">HGNC:921</a> )

**Other Names**

Galactosylgalactosylxylosylprotein  
3-beta-glucuronosyltransferase 1,  
2.4.1.135, UDP-GlcUA:glycoprotein beta-1,  
3-glucuronyltransferase, GlcUA-T-P, B3GAT1  
([http://www.genenames.org/cgi-bin/gene\\_symbol\\_report?hgnc\\_id=921](http://www.genenames.org/cgi-bin/gene_symbol_report?hgnc_id=921)  
target="\_blank">HGNC:921</a>), GLCATP


**Dilution**

IHC~~Ready-to-use

Storage	<b>Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.</b>
---------	--

Precautions	<b>CD57 Antibody is for research use only and not for use in diagnostic or therapeutic procedures.</b>
-------------	--

**CD57 Antibody - Protein Information**



Immunohistochemical analysis of paraffin-embedded human tonsil tissue using AD80190 performed on the Abcarta® FAIP-30 Fully automated IHC platform. Tissue was fixed with formaldehyde at room temperature, antigen retrieval was by heat mediation with a Citrate buffer (pH6.0). Samples were incubated with primary antibody (Ready-to-use) for 15 min at room temperature. AmpSee™ Detection Systems (Abcepta:AR005) was used as the secondary antibody.

**Name** B3GAT1 ([HGNC:921](#))**Synonyms****Function****GLCATP**

Involved in the biosynthesis of L2/HNK-1 carbohydrate epitope on glycoproteins. Can also play a role in glycosaminoglycan biosynthesis. Substrates include asialo-orosomucoid (ASOR), asialo-fetuin, and asialo-neural cell adhesion molecule. Requires sphingomyelin for activity: stearyl-sphingomyelin was the most effective, followed by palmitoyl-sphingomyelin and lignoceroyl-sphingomyelin. Activity was demonstrated only for sphingomyelin with a saturated fatty acid and not for that with an unsaturated fatty acid, regardless of the length of the acyl group.

**Cellular Location** Isoform 1: Golgi apparatus membrane {ECO:0000250|UniProtKB:O35789}; Single-pass type II membrane protein {ECO:0000250|UniProtKB:O35789}. Secreted {ECO:0000250|UniProtKB:O35789}

**Tissue Location** Mainly expressed in the brain.

## **CD57 Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)