

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM537]
Catalog # AH10958

Specification

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - Product Information

Application ,14,3,4,
Host Mouse
Clonality Monoclonal
Isotype Mouse / IgG1
Calculated MW Depends on the
target KDa

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - Additional Information

Format

200ug/ml of Ab purified from Bioreactor Concentrate by Protein A/G. Prepared in 10mM PBS with 0.05% BSA & 0.05% azide. Also available WITHOUT BSA & azide at 1.0mg/ml.

Storage

Store at 2 to 8°C.Antibody is stable for 24 months.

Precautions


Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - Protein Information

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence




Formalin-fixed, paraffin-embedded Mouse Small Intestine stained with BrdU Monoclonal Antibody (SPM537).

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - Background

It reacts with Bromodeoxyuridine (BrdU) in single stranded DNA (produced by partial denaturation of double stranded DNA), BrdU coupled to a protein carrier, as well as free BrdU. BrdU is a thymidine analog, incorporated into cell nuclei during DNA synthesis prior to mitosis. Antibody to BrdU is helpful in detecting S-phase cells, providing useful information on the aggressiveness of tumors.

Bromodeoxyuridine (BrdU) Antibody - With BSA and Azide - References

Welberg JW et. al. Journal of Clinical Pathology, 1990, 43(6):453-6. | Williams LS et. al. Cytometry, 1990, 11(4):490-7





- ImmunoprecipitationFlow CytometyCell Culture