

# FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide

Mouse Monoclonal Antibody [Clone SPM107]
Catalog # AH11247

## **Specification**

FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide - Product Information

Application ,2,3,4,
Primary Accession P01225
Other Accession 2488, 36975
Reactivity Human
Host Mouse
Clonality Monoclonal
Isotype Mouse / IgG1,

kappa

Calculated MW 22kDa KDa

FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide - Additional Information

### **Gene ID 2488**

## **Other Names**

Follitropin subunit beta, Follicle-stimulating hormone beta subunit, FSH-B, FSH-beta, Follitropin beta chain, FSHB

#### Storage

Store at 2 to 8°C.Antibody is stable for 24 months.

#### **Precautions**


FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide - Protein Information

### Name FSHB

#### **Function**

Together with the alpha chain CGA constitutes follitropin, the follicle-stimulating hormone, and provides its biological specificity to the hormone heterodimer. Binds FSHR, a G




Formalin-fixed, paraffin-embedded human Pituitary stained with FSH-beta Monoclonal Antibody (SPM107).

## FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide - Background

This MAb reacts with a protein of 22kDa, identified as beta sub-unit of FSH. It does not cross react with the alpha sub-unit. Follicle stimulating hormone (FSH) is a hormone synthesized and secreted by gonadotrophs in the anterior pituitary gland. In the ovary, FSH stimulates the growth of immature Graafian follicles to maturation. As the follicle grows, it releases inhibin, which deactivates the FSH production. In men, FSH enhances the production of androgen-binding protein by the Sertoli cells of the testis and is critical for spermatogenesis. FSH and LH act synergistically in reproduction. FSH is a useful marker in the classification of pituitary tumors and the study of pituitary disease.

## FSH-beta (Follicle Stimulating Hormone-beta) Antibody - With BSA and Azide - References

Schmidt M et al. Pathol Res Pract. 2001;197(10):663-9 | La Rosa S et al. Virchows Arch. 2000 Sep;437(3):264-9 | Zheng W et al.





Tel: 858.875.1900 Fax: 858.622.0609

protein-coupled receptor, on target cells to activate downstream signaling pathways (PubMed:<a href="http://www.uniprot.org/c itations/2494176" target="\_blank">2494176</a>, PubMed:<a href="http://www.uniprot.org/ci tations/24692546" target=" blank">24692546</a>). Follitropin is involved in follicle development and spermatogenesis in reproductive organs (PubMed:<a href="http ://www.uniprot.org/citations/407105" target=" blank">407105</a>, PubMed:<a href="http://www.uniprot.org/citations/8220 432" target=" blank">8220432</a>).

**Cellular Location** 

Secreted. Note=Efficient secretion requires dimerization with CGA

**FSH-beta** (Follicle Stimulating Hormone-beta) Antibody - With BSA and **Azide - Protocols** 

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Gynecol Oncol. 2000 jan;76(1):80-8 |