

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide
Mouse Monoclonal Antibody [Clone TNFS7/1026]
Catalog # AH12802**Specification****CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - Product Information**

Application **,13,3,4,10,5,**
Primary Accession **[P32970](#)**
Other Accession **[970](#), [501497](#),
[715224](#)**
Reactivity **Human**
Host **Mouse**
Clonality **Monoclonal**
Isotype **Mouse / IgG1,
kappa**
Calculated MW **30kDa KDa**

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - Additional Information**Gene ID 970****Other Names**

CD70 antigen, CD27 ligand, CD27-L, Tumor necrosis factor ligand superfamily member 7, CD70, CD70L, CD27LG, TNFSF7

Storage

Store at 2 to 8°C. Antibody is stable for 24 months.

Precautions

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide is for research use only and not for use in diagnostic or therapeutic procedures.

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - Protein Information**Name CD70****Synonyms CD27L, CD27LG, TNFSF7****Function**

Cytokine which is the ligand for CD27. The

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - Background

It recognizes a protein of 30kDa, identified as CD70. It is a cytokine that belongs to the tumor necrosis factor (TNF) ligand family. This cytokine is a ligand for TNFRSF27/CD27. It is a surface antigen on activated, but not on resting, T- and B-lymphocytes. It induces proliferation of co-stimulated T cells, enhances the generation of cytolytic T cells, and contributes to T cell activation. This cytokine is also reported to play a role in regulating B-cell activation, cytotoxic function of natural killer cells, and immunoglobulin synthesis.

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - References

Lens, S.M., et al. 1995. CD27-CD70 interaction: unravelling its implication in normal and neoplastic B cell growth. Leuk. Lymphoma 18: 51-59

CD70-CD27 pathway plays an important role in the generation and maintenance of T cell immunity, in particular during antiviral responses. Upon CD27 binding, induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Cellular Location

Membrane; Single- pass type II membrane protein

CD70 / TNFSF7 (Activated T- & B-Lymphocyte Marker) Antibody - With BSA and Azide - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)