

VISTA Antibody [8E11]
Catalog # ASC12162**Specification****VISTA Antibody [8E11] - Product Information**

Application	IHC-P, IF, ICC, E
Primary Accession	Q9H7M9
Other Accession	NP_071436
Host	Mouse
Clonality	Monoclonal
Isotype	IgG1
Calculated MW	Predicted: 31 kDa
	Observed: 47 kDa
	KDa

VISTA Antibody [8E11] - Additional Information

Gene ID	64115
Alias Symbol	VSIR
Other Names	
VISTA Antibody: VISTA molecule, VSIR, B7-H5, B7H5, GI24, PP2135, SISP1, DD1alpha, VISTA, C10orf54, chromosome 10 open reading frame 54, PD-1H, V-set immunoregulatory receptor, V-Type Immunoglobulin Domain-Containing Suppressor Of T-Cell Activation, Chromosome 10 Open Reading Frame 54	

Reconstitution & Storage

VISTA antibody can be stored at 4°C for three months and -20°C, stable for up to one year. As with all antibodies care should be taken to avoid repeated freeze thaw cycles. Antibodies should not be exposed to prolonged high temperatures.

Precautions

VISTA Antibody [8E11] is for research use only and not for use in diagnostic or therapeutic procedures.

VISTA Antibody [8E11] - Protein Information

Name VSIR ([HGNC:30085](#))

Function

Immunoregulatory receptor which inhibits

VISTA Antibody [8E11] - Background

VISTA Antibody: VISTA/B7-H5/platelet receptor Gi24 is a single-pass type I membrane protein located at the cell surface. It is an immunoregulatory receptor which can inhibit T-cell response and may promote differentiation of embryonic stem cells, by inhibiting the BMP4 signaling pathway. The protein can be cleaved by MMP14, and stimulate MMP14-mediated MMP2 activation.

VISTA Antibody [8E11] - References

Mayya V., et al . Quantitative phosphoproteomic analysis of T cell receptor signaling reveals system-wide modulation of protein-protein interactions. 2009, Sci. Signal. 2:RA46-RA46.Sakr M.A., et al., GI24 enhances tumor invasiveness by regulating cell surface membrane-type 1 matrix metalloproteinase. 2010, Cancer Sci. 101:2368-2374.

the T-cell response (PubMed:24691993). May promote differentiation of embryonic stem cells, by inhibiting BMP4 signaling (By similarity). May stimulate MMP14- mediated MMP2 activation (PubMed:20666777).

Cellular Location

Cell membrane; Single-pass type I membrane protein

Tissue Location

Expressed in spleen. Detected on a number of myeloid cells including CD11b monocytes, CD66b+ neutrophils, at low levels on CD4+ and CD8+ T-cells, and in a subset of NK cells. Not detected on B cells (at protein level). Expressed at high levels in placenta, spleen, plasma blood leukocytes, and lung. Expressed at moderate levels in lymph node, bone marrow, fat, uterus, and trachea Has low expression levels in other tissues

VISTA Antibody [8E11] - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)