

website:www.affbiotech.com order order@affbiotech.com

TFAM Ab

Cat.#: AF0531 Concn.: 1mg/ml Mol.Wt.: 29kDa Size: 100ul.200ul Source: Rabbit Clonality: Polyclonal

IHC 1:50-1:200 IF/ICC 1:100-1:500. WB 1:500-1:2000 Application:

Reactivity: Human, Mouse

Purification: The antiserum was purified by peptide affinity

chromatography using SulfoLink™ Coupling Resin (Thermo

Fisher Scientific).

Specificity: TFAM Ab detects endogenous levels of TFAM.

Immunogen: A synthesized peptide derived from human TFAM.

Uniprot: 000059

Description: This gene encodes a mitochondrial transcription factor that

is a key activator of mitochondrial transcription as well as a participant in mitochondrial genome replication. Studies in mice have demonstrated that this gene product is required to regulate the mitochondrial genome copy number and is essential for embryonic development. A mouse model for Kearns-Sayre syndrome was produced when expression of this gene was eliminated by targeted disruption in heart and

muscle cells.

Subcellular Location: Mitochondrion.

Binds DNA via its HMG boxes. When bound to the Similarity:

> mitochondrial light strand promoter, bends DNA into a Uturn shape, each HMG box bending the DNA by 90 degrees.

Storage Condition and

Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20


°C.Stable for 12 months from date of receipt.



Western blot analysis of extracts from mouse brain, using TFAM Ab. Lane 1 was treated with the antigen-specific peptide.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



AF0531 at 1/100 staining human brain tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.