


Phospho-JAK2 (Tyr570) Ab

Cat.#: AF3025
Size: 100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 125kDa
Clonality: Polyclonal

Application:	WB 1:500-1:2000, IF/ICC 1:100-1:500
Reactivity:	Human,Mouse,Rat
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Phospho-JAK2 (Tyr570) Ab detects endogenous levels of JAK2 only when phosphorylated at Tyrosine 570.
Immunogen:	A synthesized peptide derived from human JAK2 around the phosphorylation site of Tyrosine 570.
Uniprot:	O60674
Description:	This gene product is a protein tyrosine kinase involved in a specific subset of cytokine receptor signaling pathways. It has been found to be constitutively associated with the prolactin receptor and is required for responses to gamma interferon.
Subcellular Location:	Endomembrane system. Nucleus.
Tissue Specificity:	Ubiquitously expressed throughout most tissues.
Similarity:	Possesses 2 protein kinase domains. The second one probably contains the catalytic domain, while the presence of slight differences suggest a different role for protein kinase 1 (By similarity).Belongs to the protein kinase superfamily. Tyr protein kinase family. JAK subfamily.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-JAK2 (Tyr570) Ab expression in etoposide treated 293 cells lysates. The lane on the right is treated with the antigen-specific peptide.



Western blot analysis of JAK2 phosphorylation expression in etoposide treated 293 whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3025 staining K562 by IF/ICC. The sample were fixed with PFA and permeabilized in 0.1% Triton X-100, then blocked in 10% serum for 45 minutes at 25°C. The primary Ab was diluted at 1/200 and incubated with the sample for 1 hour at 37°C. An Alexa Fluor 594 conjugated goat anti-rabbit IgG (H+L) Ab, diluted at 1/600, was used as the secondary Ab.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.