


Phospho-iNOS (Tyr151) Ab

Cat.#: AF3270
Size: 100ul,200ul


Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: kDa
Clonality: Polyclonal

Application:	IHC 1:50-1:200 WB 1:500
Reactivity:	Human,Mouse,Rat
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Phospho-iNOS (Tyr151) Ab detects endogenous levels of iNOS only when phosphorylated at Tyrosine 151.
Immunogen:	A synthesized peptide derived from human iNOS around the phosphorylation site of Tyrosine 151.
Uniprot:	P35228
Description:	iNOS Produces nitric oxide (NO) which is a messenger molecule with diverse functions throughout the body. In macrophages, NO mediates tumoricidal and bactericidal actions.
Tissue Specificity:	Expressed in the liver, retina, bone cells and airway epithelial cells of the lung. Not expressed in the platelets.
Similarity:	Belongs to the NOS family.
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of iNOS phosphorylation expression in whole cell lysates, The lane on the left is treated with the antigen-specific peptide.



AF3270 at 1/100 staining human colon tissue sections by IHC-P. The tissue was formaldehyde fixed and a heat mediated antigen retrieval step in citrate buffer was performed. The tissue was then blocked and incubated with the Ab for 1.5 hours at 22°C. An HRP conjugated goat anti-rabbit Ab was used as the secondary.

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.