## Phospho-TrkA (Tyr785) /TrkB (Tyr816) Ab

Cat.#: AF4390 Concn.: 1mg/ml Mol.Wt.: 87kDa Size: 50ul,100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-TrkA (Tyr785) /TrkB (Tyr816) Ab detects

endogenous levels of TrkA only when phosphorylated at

Tyr785.

Immunogen: A synthesized peptide derived from human TrkA around the

phosphorylation site of Tyr785.

Uniprot: P04629

Subcellular Location: Cell membrane. Early endosome membrane. Late endosome

membrane. Internalized to endosomes upon binding of NGF or NTF3 and further transported to the cell body via a retrograde axonal transport. Localized at cell membrane and

early endosomes before nerve growth factor (NGF) stimulation. Recruited to late endosomes after NGF stimulation. Colocalized with RAPGEF2 at late endosomes

(By similarity).

Tissue Specificity: Isoform TrkA-I is found in most non-neuronal tissues. Isoform

TrkA-II is primarily expressed in neuronal cells. TrkA-III is specifically expressed by pluripotent neural stem and neural

crest progenitors.

Similarity: The transmembrane domain mediates interaction with

KIDINS220.The extracellular domain mediates interaction with NGFR.Belongs to the protein kinase superfamily. Tyr

protein kinase family. Insulin receptor subfamily.

Storage Condition and


Buffer:

Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



## Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of Phospho-TrkA (Tyr785) /TrkB (Tyr816) in lysates of jurkat , using Phospho-TrkA (Tyr785) /TrkB (Tyr816) Ab(AF4390).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at  $4^{\circ}$ C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.