Phospho-FLT3 (Tyr842) Ab

Cat.#: AF4427 Concn.: 1mg/ml Mol.Wt.: 113kDa Size: 50ul,100ul,200ul Source: Rabbit Clonality: Polyclonal

Application: WB 1:500-1:2000

Reactivity: Human, Mouse, Rat

Purification: The Ab is from purified rabbit serum by affinity purification

via sequential chromatography on phospho- and non-

phospho-peptide affinity columns.

Specificity: Phospho-FLT3 (Tyr842) Ab detects endogenous levels of

FLT3 only when phosphorylated at Tyr842.

Immunogen: A synthesized peptide derived from human FLT3 around the

phosphorylation site of Tyr842.

Uniprot: P36888

Subcellular Location: Membrane.

Tissue Specificity: Detected in bone marrow, in hematopoietic stem cells, in

myeloid progenitor cells and in granulocyte/macrophage progenitor cells (at protein level). Detected in bone marrow, liver, thymus, spleen and lymph node, and at low levels in kidney and pancreas. Highly expressed in T-cell leukemia.

Similarity: The juxtamembrane autoregulatory region is important for

normal regulation of the kinase activity and for maintaining the kinase in an inactive state in the absence of bound ligand. Upon tyrosine phosphorylation, it mediates interaction with the SH2 domains of numerous signaling partners. In-frame internal tandem duplications (ITDs) result in constitutive activation of the kinase. The activity of the mutant kinase can be stimulated further by FLT3LG binding.Belongs to the protein kinase superfamily. Tyr protein kinase family. CSF-1/PDGF receptor subfamily.

Storage Condition and


Buffer:

Rabbit IgG in phosphate buffered saline, pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol. Store at -20

°C.Stable for 12 months from date of receipt.



Affinity Biosciences website:www.affbiotech.com order:order@affbiotech.com



Western blot analysis of Phospho-FLT3 (Tyr842) in lysates of COS7 EGF 200ng/ml 30', using Phospho-FLT3 (Tyr842) Ab(AF4427).

<code>IMPORTANT:</code> For western blot, incubate membrane with diluted primary Ab in 5% w/v milk , 1X TBS, 0.1% Tween020 at 4° C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.