

Phospho-53BP1 (Ser1618) Ab

Cat.#: AF4460
Size: 50ul,100ul,200ul

Concn.: 1mg/ml
Source: Rabbit

Mol.Wt.: 450kDa
Clonality: Polyclonal

Application:	WB 1:500-1:2000
Reactivity:	Human,Rat
Purification:	The Ab is from purified rabbit serum by affinity purification via sequential chromatography on phospho- and non-phospho-peptide affinity columns.
Specificity:	Phospho-53BP1 (Ser1618) Ab detects endogenous levels of 53BP1 only when phosphorylated at Ser1618.
Immunogen:	A synthesized peptide derived from human 53BP1 around the phosphorylation site of Ser1618.
Uniprot:	Q12888
Subcellular Location:	Nucleus. Chromosome > centromere > kinetochore. Associated with kinetochores. Both nuclear and cytoplasmic in some cells. Recruited to sites of DNA damage, such as double strand breaks. Methylation of histone H4 at 'Lys-20' is required for efficient localization to double strand breaks.
Similarity:	The Tudor-like region mediates binding to histone H4 dimethylated at 'Lys-20' (H4K20me2) (PubMed:17190600). Interaction with NUDT16L1/TIRR masks the Tudor-like domain and prevents recruitment to chromatin (PubMed:28241136).The UDR (ubiquitin-dependent recruitment) motif specifically recognizes and binds histone H2A monoubiquitinated at 'Lys-15' (H2AK15ub) (PubMed:23760478, PubMed:24703952). Phosphorylation of the UDR blocks interaction with H2AK15ub (PubMed:24703952).
Storage Condition and Buffer:	Rabbit IgG in phosphate buffered saline , pH 7.4, 150mM NaCl, 0.02% sodium azide and 50% glycerol.Store at -20 °C.Stable for 12 months from date of receipt.



Western blot analysis of Phospho-53BP1 (Ser1618) in lysates of HeLa thymidine nocodazole, using Phospho-53BP1 (Ser1618) Ab(AF4460).

IMPORTANT: For western blot, incubate membrane with diluted primary Ab in 5% w/v milk, 1X TBS, 0.1% Tween@20 at 4°C with gentle shaking, overnight.

For Research Use Only. Not for use in diagnostic and therapeutic procedures. Not for resale without express authorization.