



Mouse Monoclonal Antibody to CD10

Cataloge Number sAP-0872

Target Molecule Name: CD10

Aliases: MME; NEP; SFE; CALLA

MW: 85.5kDa

Entrez Gene ID: 4311

Descrption

This gene encodes a common acute lymphocytic leukemia antigen that is an important cell surface marker in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of

in the diagnosis of human acute lymphocytic leukemia (ALL). This protein is present on leukemic cells of pre-B phenotype, which represent 85% of cases of ALL. This protein is not restricted to leukemic cells, however, and is found on a variety of normal tissues. It is a glycoprotein that is particularly abundant in kidney, where it is present on the brush border of proximal tubules and on glomerular epithelium. The protein is a neutral endopeptidase that cleaves peptides at the amino side of hydrophobic residues and inactivates several peptide hormones including glucagon, enkephalins, substance P, neurotensin, oxytocin, and bradykinin. This gene, which encodes a 100-kD type II transmembrane glycoprotein, exists in a single copy

Immunogen Purified recombinant fragment of human CD10 (AA: 52-246) expressed in E. Coli.

Recitative Species Human;

Clone MM5E9A5;

Size and Concentration 100µg/1mg/ml

Supplied as Lyophilized Powder from 100µl of Purified antibody in PBS with 0.05% sodium azide

Reconstitution/Storages Reconstituted with 100µl sterile DI H2O, at stored at 4°C or -20°C for short or long term storage

Applications ELISA: 1 to 10000; WB: 1 to 500 - 1 to 2000; IHC: 1 to 200 - 1 to 1000

Shipping Regular FEDEX overnight shipment (ambient temperature)

Reference 1. Pathol Res Pract. 2012 May 15;208(5):281-5.; 2. J Dermatol Sci. 2013 Feb;69(2):105-13.;

Optimal dilutions should be determined by each laboratory for each application. The listed dilutions are for recommendation only and the final conditions should be optimized by the ender users! This product is sold for Research Use Only