

## Rabbit Anti-Rat Histamine pAb

<b>Catalog No.</b>	PS242	<b>Quantity:</b>	50 µl
<b>Specificity:</b>	Positive control: Cryostat sections of carbodiimide fixed human skin or freeze-dried paraffin sections (vapour fixed in DEPC) of rat stomach (see ref.). Absorption with 10-100 µg histamine per ml diluted antiserum abolishes the staining, while noradrenaline, 5-HT, VIP, glucagon and histidine do not.		
<b>Host:</b>	Rabbit		
<b>Immunogen:</b>	Histamine conjugated to HSA. Tested on rat tissues.		
<b>Formulation:</b>	50 µl undiluted, freeze-dried antiserum. Dissolve the antiserum in 50-100 µL distilled water, and dilute further in 0.1M PBS with 1% BSA and 0.1% NaN <sub>3</sub> .		
<b>Applications:</b>	Works on: Paraffin and frozen sections. 1:1500-1:2000 using PAP, or 1:300-1:600 using FITC with over night incubation at +4°C.		
<b>Storage &amp; Stability:</b>	Freeze-dried at 2-8°C until expiration date. Reconstituted in small aliquots at -20°C.		

**NOT FOR HUMAN USE. FOR RESEARCH ONLY. NOT FOR DIAGNOSTIC OR THERAPEUTIC USE.**