



Recombinant SARS-CoV-2 Spike S1 (K417N, E484K, N501Y, D614G) Protein [His] (DAGC489)

This product is for research use only and is not intended for diagnostic use.

PRODUCT INFORMATION

Product Overview	A DNA sequence encoding the SARS-CoV-2 (2019-nCoV) Spike S1(K417N, E484K, N501Y, D614G)-His Recombinant Protein (YP_009724390.1) (Met1-Arg685(K417N, E484K, N501Y, D614G)) was expressed with a polyhistidine tag at the C-terminus.
Nature	Recombinant
Expression System	HEK293 Cells
Species	SARS-CoV-2
Purity	> 90 % as determined by SDS-PAGE
Conjugate	His
Applications	ELISA
Molecular Weight	The recombinant SARS-CoV-2 (2019-nCoV) Spike S1(K417N, E484K, N501Y, D614G)-His Recombinant Protein consists of 681 amino acids and predicts a molecular mass of 76.4 kDa. As a result of glycosylation, it migrates as an approximately 114.7 kDa band in SDS-PAGE under reducing conditions.
Procedure	None
Format	Lyophilized
Size	100ug
Buffer	Lyophilized from sterile PBS, pH 7.4. Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization.
Preservative	None
Storage	Store it under sterile conditions at -20°C to -80°C. It is recommended that the protein be aliquoted for optimal storage. Avoid repeated freeze-thaw cycles.

BACKGROUND

Introduction

In South Africa, another variant of SARS-CoV-2 (known as 20H/501Y.V2 or B.1.351) emerged independently of B.1.1.7. This variant has multiple mutations in the spike protein, including K417T, E484K, N501Y. Unlike the B.1.1.7 lineage detected in the UK this variant does not contain the deletion at 69/70. The spike (S) glycoprotein of coronaviruses contains protrusions that will only bind to certain receptors on the host cell: they are essential for both host specificity and viral infectivity. The term 'peplomer' is typically used to refer to a grouping of heterologous proteins on the virus surface that function together.

The spike (S) glycoprotein of coronaviruses is known to be essential in the binding of the virus to the host cell at the advent of the infection process. Most notable is severe acute respiratory syndrome (SARS). The severe acute respiratory syndrome-coronavirus (SARS-CoV) spike (S) glycoprotein alone can mediate the membrane fusion required for virus entry and cell fusion. It is also a major immunogen and a target for entry inhibitors. The SARS-CoV spike (S) protein is composed of two subunits; the S1 subunit contains a receptor-binding domain that engages with the host cell receptor angiotensin-converting enzyme 2 and the S2 subunit mediates fusion between the viral and host cell membranes. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity, during infection with SARS-CoV.

Keywords

SARS-CoV-2 Spike S1; SARS-CoV-2; SARS-CoV-2 S1; SARS-CoV-2 Spike
