





## CXCR2 Antibody, FITC conjugated

| <b>Product Code</b> | CSB-PA011673HC01HU   |
|---------------------|--|
| Storage             | Upon receipt, store at -20°C or -80°C. Avoid repeated freeze.  |
| Uniprot No.         | P25025   |
| Immunogen           | Recombinant Human C-X-C chemokine receptor type 2 protein (1-40AA)   |
| Raised In           | Rabbit   |
| Species Reactivity  | Human  |
| Tested Applications | ELISA  |
| Relevance           | Receptor for interleukin-8 which is a powerful neutrophil chemotactic factor. Binding of IL-8 to the receptor causes activation of neutrophils. This response is mediated via a G-protein that activates a phosphatidylinositol-calcium second messenger system. Binds to IL-8 with high affinity. Also binds with high affinity to CXCL3, GRO/MGSA and NAP-2. |
| Form                | Liquid   |
| Conjugate           | FITC   |
| Storage Buffer      | Preservative: 0.03% Proclin 300<br>Constituents: 50% Glycerol, 0.01M PBS, PH 7.4   |
| Purification Method | >95%, Protein G purified   |
| Isotype             | IgG  |
| Clonality           | Polyclonal   |
| Alias               | C-X-C chemokine receptor type 2 (CXC-R2) (CXCR-2) (CDw128b) (GRO/MGSA receptor) (High affinity interleukin-8 receptor B) (IL-8R B) (IL-8 receptor type 2) (CD antigen CD182), CXCR2, IL8RB   |
| Species             | Homo sapiens (Human)   |
| Research Area       | Signal Transduction  |
| Target Names        | CXCR2  |