

IL-8Rβ (phospho Ser347) Polyclonal Antibody

| Swiss-Prot No.: | P25025 |
|--------------------------|--|
| Molecular Weight: | 40.759 |
| Applications: | WB,IHC,IF,ELISA |
| Reactivity: | Human,Mouse |
| Purification: | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Immunogen: | Synthesized peptide derived from human IL-8R β around the phosphorylation site of S347. |
| Specificity/Sensitivity: | Phospho-IL-8R β (S347) Polyclonal Antibody detects endogenous levels of IL-8R β protein only when phosphorylated at S347. |
| Other Names: | CXCR2; IL8RB; C-X-C chemokine receptor type 2; CXC-R2; CXCR-2; CDw128b; GRO/MGSA receptor; High affinity interleukin-8 receptor B; IL-8R B; IL-8 receptor type 2; CD antigen CD182 |
| Storage/Stability: | -20°C/1 year |
| Source: | Rabbit |
| Form of Antibody: | Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. |

| Protein Concentration: | 1 mg/ml |
|------------------------|--|
| | 3T3 178 100 70 55 40 25 15 PMA P-IL-8Rβ (S347) |
| | Western Blot analysis of 3T3 cells using Phospho-IL-8R β (S347) Polyclonal Antibody |
| Gene Name: | CXCR2 |
| Dilution: | Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications. |