




## IL-8 Polyclonal Antibody

<b>Swiss-Prot No.:</b>	P10145
<b>Molecular Weight:</b>	11.098
<b>Applications:</b>	WB,IHC,ELISA
<b>Reactivity:</b>	Human
<b>Purification:</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Immunogen:</b>	Synthesized peptide derived from the Internal region of human IL-8.
<b>Specificity/Sensitivity:</b>	IL-8 Polyclonal Antibody detects endogenous levels of IL-8 protein.
<b>Other Names:</b>	IL8; CXCL8; Interleukin-8; IL-8; C-X-C motif chemokine 8; Emotakin; Granulocyte chemotactic protein 1; GCP-1; Monocyte-derived neutrophil chemotactic factor; MDNCF; Monocyte-derived neutrophil-activating peptide; MONAP; Neutrophil-activati
<b>Storage/Stability:</b>	-20°C/1 year
<b>Source:</b>	Rabbit
<b>Form of Antibody:</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.

**For Research Use Only**

<b>Protein Concentration:</b>	1 mg/ml
	<p style="text-align: center;">HeLa</p>  <p>(kD)</p> <p>117-</p> <p>85-</p> <p>48-</p> <p>34-</p> <p>26-</p> <p>19-</p>
	Western Blot analysis of various cells using IL-8 Polyclonal Antibody
<b>Gene Name:</b>	IL8
<b>Dilution:</b>	Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.