

DATASHEET

Version 20181206

sCD40L, Human

Cat. No.: Z02727-1

Size: 1.0 mg

Synonyms: soluble CD40 Ligand (sCD40L), Human;

Description:

CD40 ligand, CD40L (also known as CD154, TRAP or gp39), is a 261 amino acid type II transmembrane glycoprotein belonging to the TNF family, CD40L is expressed predominantly on activated CD4+ T lymphocytes, and also found in other types of cells, like NK cells, mast cells, basophils and eosinophils. Human CD40L shares 78% amino acid identity with its mouse counterpart. The receptor of CD40L is CD40, a type I transmembrane glycoprotein belonging to the TNF receptor family. CD40 is expressed on B lymphocytes, monocytes, dendritic cells and thymic epithelium. Although all monomeric, dimeric and trimeric forms of soluble CD40L can bind to CD40, the trimeric form of soluble CD40L has the most potent biological activity through oligomerization of cell surface CD40, a common feature of TNF receptor family members. CD40L mediates a range of activities on B cells including induction of activation-associated surface antigen, entry into cell cycle, isotype switching and Ig secretion and memory generation. CD40-CD40L interaction also plays important roles in monocyte activation and dendritic cell maturation.

Amino Acid Sequence:

```
00001 MQKGDQNPQI AAHVISEASS KTTSVLQWAE KGYTMSNNL
00041 VTLENGKQLT VKRQGLYYIY AQVTFCSNRE ASSQAPFIAS
00081 LWLKSPGRFE RILLRAANTH SSAKPCGQQS IHLGGVFELQ
00121 PGASVFVNVT DPSQVSHGTG FTSFGLLKL
```

Source: *E. coli*

Species: Human

Biological Activity: Fully biologically active when compared to standard. The ED₅₀ as determined by the dose-dependent stimulation of IL-8 production by human PBMC is less than 5-10 ng/ml.

Molecular Weight: Approximately 16.3 kDa, a single non-glycosylated polypeptide chain containing 149 amino acids.

Formulation: Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.0.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at ≤ -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 95 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/µg of rHusCD40L as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.