

DATASHEET
Version 20181206**IL-6, RhesusMacaque****Cat. No.:** Z02759-10**Size:** 10.0 ug**Synonyms:** Interleukin-6 (IL-6), Rhesus Macaque;**Description:**

IL-6 is a pleiotropic cytokine that plays an important role in host defense by regulating immune and inflammatory responses. Produced by T cells, monocytes, fibroblasts, endothelial cells and keratinocytes, IL-6 has diverse biological functions. It stimulates B-cell differentiation and antibody production, synergizes with IL-3 in megakaryocyte development and platelet production, induces expression of hepatic acute-phase proteins, and regulates bone metabolism. IL-6 signals through the IL-6 receptor system that consists of two chains, IL-6R^a and gp130.

Amino Acid Sequence:

```
00001 MAPVLPGEDS KNVAAPHSQP LTSSERIDKH IRYILDGISA
00041 LRKETCNRSN MCESSKEALA ENNLNLPKMA EKDGCFQSGF
00081 NEDTCLVKII TGLLEFEVYL EYLQNRFESE EQARAVQMS
00121 TKVLIQFLQK KAKNLDAITT PEPTNASLL TKLQAQNWQL
00161 QDMTTHLILR SFKEFLQSNL RALRQM
```

Source: *E. coli***Species:** RhesusMacaque

Biological Activity: Fully biologically active when compared to standard. The ED₅₀ as determined by a cell proliferation assay using IL-6-dependent murine 7TD1 cells is less than 0.1 ng/ml, corresponding to a specific activity of $> 1.0 \times 10^7$ IU/mg.

Molecular Weight: Approximately 21.1 kDa, a single non-glycosylated polypeptide chain containing 186 amino acids.

Formulation: Lyophilized from a 0.2 µm filtered concentrated solution in PBS, pH 7.4.

Appearance: Sterile Filtered White lyophilized (freeze-dried) powder.

Reconstitution: We recommend that this vial be briefly centrifuged prior to opening to bring the contents to the bottom. Reconstitute in sterile distilled water or aqueous buffer containing 0.1 % BSA to a concentration of 0.1-1.0 mg/mL. Stock solutions should be apportioned into working aliquots and stored at < -20 °C. Further dilutions should be made in appropriate buffered solutions.

Purity: > 97 % by SDS-PAGE and HPLC analyses.

Endotoxin Level: Less than 1 EU/µg of rRhIL-6 as determined by LAL method.

Storage: This lyophilized preparation is stable at 2-8 °C, but should be kept at -20 °C for long term storage, preferably desiccated. Upon reconstitution, the preparation is stable for up to one week at 2-8 °C. For maximal stability, apportion the reconstituted preparation into working aliquots and store at -20 °C to -70 °C. Avoid repeated freeze/thaw cycles.