

Human B7-H5/Gi24/VISTA Protein

Cat. No. BH7-HM275

Description

Source	Recombinant Human B7-H5/Gi24/VISTA Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Phe33-Ala194.
Accession	Q9H7M9
Molecular Weight	The protein has a predicted MW of 44.9 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

B7-H5, also known as VISTA, B7H5, Dies1, SISP1 and C10orf54, is a 55-65 kDa member of the Ig superfamily. It is a transmembrane molecule expressed in bone, on embryonic stem cells (ESCs), and on tumor cell surfaces.


Assay Data

Tris-Bis PAGE



Human B7-H5 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human B7-H5 is greater than 95% as determined by SEC-HPLC

Assay Data


ELISA Data

Human B7-H5, hFc Tag ELISA
0.05µg Human B7-H5, hFc Tag Per Well



Immobilized Human B7-H5, hFc Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-B7-H5 Antibody, hFc Tag with the EC50 of 8.0ng/ml determined by ELISA (QC Test).

SPR Data



Human B7-H5, hFc Tag captured on CM5 Chip via Protein A can bind Human IGSF11, His Tag with an affinity constant of 14.44 µM as determined in SPR assay (Biacore T200).