

Description

Source	Recombinant Biotinylated Human B7-H3 (4Ig)/B7-H3b Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Gly27-Thr461.
Accession	Q5ZPR3-1
Molecular Weight	The protein has a predicted MW of 49.5 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

B7-H3, a member of the B7 family of immunomodulatory molecules, is overexpressed in a wide range of solid cancers. B7-H3 binds to activated T cells via an as yet unidentified receptor. In assays using sub-optimal amount so anti-CD3 stimulation, 2IgB7H3 enhances T cell proliferation, T cell interferon-gamma (IFN-gamma) production, and cytotoxic T cells induction.

Assay Data


Tris-Bis PAGE



Biotinylated Human B7-H3 (4Ig) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Assay Data



The purity of Biotinylated Human B7-H3 (4Ig) is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human B7-H3 (4Ig), His Tag ELISA0.05 μ g Anti-B7-H3 Antibody, hFc Tag Per Well

Immobilized Anti-B7-H3 Antibody, hFc Tag at 0.5 μ g/ml (100 μ l/well) on the plate. Dose response curve for Biotinylated Human B7-H3 (4Ig) , His Tag with the EC50 of 7.7ng/ml determined by ELISA.