

Human CD3E/CD3 epsilon 1-27 peptide Protein



Cat. No. CD3-HM2ED


Description	
Source	Recombinant Human CD3E/CD3 epsilon 1-27 peptide Protein is expressed from HEK293 with hFc tag and Avi tag at the C-Terminus. It contains Asp23-Thr48.
Accession	P07766
Molecular Weight	The protein has a predicted MW of 31.3 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
CD3E, is a single-pass type I membrane protein. CD3 (cluster of differentiation 3) T cell co-receptor helps to activate both the cytotoxic T cell (CD8 naive T cells) and also T helper cells (CD4 naive T cells). It consists of a protein complex and is composed of four distinct chains. In mammals, the complex contains a CD3γ chain, a CD3δ chain, and two CD3ε chains.	

Assay Data

Tris-Bis PAGE



Human CD3E 1-27 peptide on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Human CD3E/CD3 epsilon 1-27 peptide Protein

Cat. No. CD3-HM2ED




Assay Data



The purity of Human CD3E 1-27 peptide is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human CD3E 1-27 peptide, hFc Tag ELISA
0.1µg Human CD3E 1-27 peptide, hFc Tag Per Well



Immobilized Human CD3E 1-27 peptide, hFc Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-CD3 Antibody, mFc Tag with the EC50 of 5.2ng/ml determined by ELISA.