Mouse CD27 Ligand/CD70 Protein

Cat. No. CD7-MM270



Description	
Source	Recombinant Mouse CD27 Ligand/CD70 Protein is expressed from HEK293 with hFc tag at the N-Terminus.
	It contains Ser45-Pro195.
Accession	O55237
Molecular Weight	The protein has a predicted MW of 43.41 kDa. Due to glycosylation, the protein migrates to 55-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


· · · · · · · · · · · · · · · · · · ·	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 6.2). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Assay Data

Tris-Bis PAGE




Mouse CD27 Ligand on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS

Assay Data




The purity of Mouse CD27 Ligand is greater than 95% as determined by SEC-HPLC.

ELISA Data

Mouse CD27 Ligand, hFc Tag ELISA

0.1μg Mouse CD27, mFc Tag Per Well



Immobilized Mouse CD27, mFc Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Mouse CD27 Ligand, hFc Tag with the EC50 of 35.5ng/ml determined by ELISA.