
Biotinylated Human CD27 Ligand/CD70 Trimer Protein (Primary Amine Labeling)

Dissolve the lyophilized protein in distilled water.





Description	
Source	Recombinant Biotinylated Human CD27 Ligand/CD70 Trimer Protein (Primary Amine Labeling) is expressed from HEK293 with His tag at the N-Terminus.
	It contains Leu50-Pro193.
Accession	P32970-1
Molecular Weight	The protein has a predicted MW of 51.8 kDa. Due to glycosylation, the protein migrates to 55-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	Storage
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended.

Storage

-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.


Background

Reconstitution

CD70, also named CD27 ligand (CD27L), is a type II transmembrane glycoprotein belonging to the TNF superfamily (TNFSF) and has been designated TNFSF7. CD70 is a cytokine that binds to CD27. Plays a role in T-cell activation. Induces the proliferation of costimulated T-cells and enhances the generation of cytolytic T-cells.

Assay Data

Tris-Bis PAGE




Biotinylated Human CD27 Ligand (Trimer) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




Assay Data



The purity of Biotinylated Human CD27 Ligand (Trimer) is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated Human CD27 Ligand (Trimer), HisTag ELISA 0.1µg Human CD27, hFc Tag Per Well



Immobilized Human CD27, hFc Tag at $1\mu g/ml$ ($100\mu l/Well$) on the plate. Dose response curve for Biotinylated Human CD27 Ligand (Trimer) , His Tag with the EC50 of 9.2ng/ml determined by ELISA.