
# Biotinylated SARS-COV-2 Spike RBD Protein





Description	
Source	Recombinant Biotinylated SARS-COV-2 Spike RBD Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Arg319-Asn532.
Accession	QHD43416.1
Molecular Weight	The protein has a predicted MW of 27 kDa. Due to glycosylation, the protein migrates to 36-40 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and	4 Storago

### Formulation and Storage


Formulation	lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

**Background** 

The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

## **Assay Data**

### **Tris-Bis PAGE**




Biotinylated SARS-COV-2 Spike RBD on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

# KAGTUS


## **Assay Data**



The purity of Biotinylated SARS-COV-2 Spike RBD is greater than 95% as determined by SEC-HPLC.

### **ELISA Data**

# Biotinylated SARS-COV-2 Spike RBD, His Tag ELISA 0.1µg Biotinylated SARS-COV-2 Spike RBD, His Tag Per Well



Immobilized Biotinylated SARS-COV-2 Spike RBD, His Tag at 1µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Human ACE2, hFc Tag with the EC50 of 7.7ng/ml determined by ELISA.