

Mouse ICOS/CD278 Protein



Cat. No. ICO-MM201


Description	
Source	Recombinant Mouse ICOS/CD278 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains Glu21-Leu144.
Accession	Q9WVS0-1
Molecular Weight	The protein has a predicted MW of 40.8 kDa. Due to glycosylation, the protein migrates to 65-80 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Inducible co-stimulator (ICOS), also called AILIM (activation-inducible lymphocyte immunomediatory molecule) and CRP-1 (CD28-related protein-1), is a member of the growing CD28 family of immune costimulatory receptors. Other family members are CD28, CTLA-4 and PD-1. Human ICOS is a homodimeric type I transmembrane protein consisting of 199 amino acids (aa) with a putative 20 aa signal sequence, a 121 aa extracellular domain, a 23 aa transmembrane region, and a 35 aa cytoplasmic domain.	

Assay Data

Tris-Bis PAGE



Mouse ICOS on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Mouse ICOS/CD278 Protein

Cat. No. ICO-MM201




Assay Data



The purity of Mouse ICOS is greater than 95% as determined by SEC-HPLC.

SPR Data



Mouse ICOS, hFc Tag captured on CM5 Chip via Protein A can bind Mouse B7-H2, His Tag with an affinity constant of 0.329 μM as determined in SPR assay (Biacore T200).