SARS-Cov-2 Spike RBD (B.1.640.2/IHU) Protein





| Description | |
|---------------------|---|
| Source | Recombinant SARS-Cov-2 Spike RBD(B.1.640.2/IHU) Protein is expressed from HEK293 with His tag at the C-Terminus. |
| | It contains Arg319-Phe541(R346S, N394S, Y449N, E484K, F490S, N501Y). |
| Accession | QHD43416.1 |
| Molecular Weight | The protein has a predicted MW of 25.76 kDa. Due to glycosylation, the protein migrates to 35-40 kDa based on Tris-Bis PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Tris-Bis PAGE |
| | > 95% as determined by HPLC |
| Formulation and | d Storage |


| Formulation | lyophilization. |
|----------------|--|
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Assay Data

Tris-Bis PAGE




SARS-Cov-2 Spike RBD (B.1.640.2/IHU) on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. IHU-VM1RD

KAGTUS


Assay Data



The purity of SARS-Cov-2 Spike RBD (B.1.640.2/IHU) is greater than 95% as determined by SEC-HPLC.

ELISA Data

SARS-Cov-2 Spike RBD(B.1.640.2/IHU), His Tag ELISA 0.05µg SARS-Cov-2 Spike RBD(B.1.640.2/IHU), His Tag Per Well



Immobilized SARS-Cov-2 Spike RBD (B.1.640.2/IHU) , His Tag at $0.5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Human ACE2, hFc Tag with the EC50 of 41.3ng/ml determined by ELISA.