

Biotinylated Human IL-17A&F Protein



Cat. No. ILF-HM418B


Description	
Source	Recombinant Biotinylated Human IL-17A&F Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Gly24-Ala155(IL-17A)&Arg31-Gln163(IL-17F).
Accession	Q16552-1(IL-17A)&Q96PD4(IL-17F)
Molecular Weight	The protein has a predicted MW of 15.1 kDa (IL-17A)&17.5 kDa (IL-17F). Due to glycosylation, the protein migrates to 16 kDa&27 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background	
Interleukin17A (IL17A), also known as CTLA8, is a 1520 kDa glycosylated cytokine that plays an important role in antimicrobial and chronic inflammation. The six IL17 cytokines (IL17AF) are encoded by separate genes but adopt a conserved cystine knot fold.IL-17A is a ligand for IL17RA and IL17RC. The heterodimer formed by IL17A and IL17F is a ligand for the heterodimeric complex formed by IL17RA and IL17RC. Involved in inducing stromal cells to produce proinflammatory and hematopoietic cytokines.	

Assay Data


Tris-Bis PAGE



Biotinylated Human IL-17A&F on Tris-Bis PAGE under reduced conditions. The purity is greater than 95%.

ELISA Data

Biotinylated Human IL-17A&F, His Tag ELISA  
0.2µg Human IL-17R alpha, hFc Tag Per Well



Immobilized Human IL-17R alpha, hFc Tag at 2µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Human IL-17A&F, His Tag with the EC50 of 0.22µg/ml determined by ELISA.