

Biotinylated SARS Spike RBD Protein

Cat. No. SAR-VM4BDB



Description

Source	Recombinant Biotinylated SARS RBD Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus.
	It contains Arg306-Phe527.
Accession	P59594
Molecular Weight	The protein has a predicted MW of 27.9 kDa. Due to glycosylation, the protein migrates to 36-46 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The spike protein (S) of coronavirus (CoV) attaches the virus to its cellular receptor, angiotensin-converting enzyme 2 (ACE2). A defined receptor-binding domain (RBD) on S mediates this interaction. The S protein plays key parts in the induction of neutralizing-antibody and T-cell responses, as well as protective immunity.

Assay Data

Tris-Bis PAGE



Biotinylated SARS RBD on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated SARS Spike RBD Protein

Cat. No. SAR-VM4BDB



Assay Data




The purity of Biotinylated SARS RBD is greater than 95% as determined by SEC-HPLC.

ELISA Data

Biotinylated SARS Spike RBD, His Tag ELISA

0.5µg Human ACE2, hFc Tag Per Well



Immobilized Human ACE2, hFc Tag at 5µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated SARS Spike RBD, His Tag with the EC50 of 0.19µg/ml determined by ELISA.