Human Tim-3/HAVCR2 Protein

Cat. No. TIM-HM131



| Description | |
|---------------------|--|
| Source | Recombinant Human Tim-3/HAVCR2 Protein is expressed from HEK293 with His tag at the C-Terminus. |
| | It contains Ser22-Arg200. |
| Accession | Q8TDQ0-1 |
| Molecular Weight | The protein has a predicted MW of 20.7 kDa, Due to glycosylation, the protein migrates to 50-60 kDa based on Tris-Bis PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Tris-Bis PAGE |
| | > 95% as determined by HPLC |
| | |

Formulation and Storage


| romulation and Storage | |
|------------------------|--|
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μ g/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Hepatitis A virus cellular receptor 2 (HAVCR2), also known as T-cell immunoglobulin and mucin-domain containing-3 (TIM-3), is a protein that in humans is encoded by the HAVCR2 gene.TIM3 is an immune checkpoint and together with other inhibitory receptors including programmed cell death protein 1 (PD-1) and lymphocyte activation gene 3 protein (LAG3) mediate the CD8 T-cell exhaustion. TIM3 has also been shown as a CD4 Th1-specific cell surface protein that regulates macrophage activation and enhances the severity of experimental autoimmune encephalomyelitis in mice.

Assay Data

Tris-Bis PAGE




Human TIM-3 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Human TIM-3 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human Tim-3, His Tag ELISA 0.05µg Human Tim-3, His Tag Per Well



Immobilized Human TIM-3, His Tag at 0.5µg/ml (100µl/well) on the plate. Dose response curve for Anti-TIM-3 Antibody, hFc Tag with the EC50 of 4.5ng/ml determined by ELISA.