

Human CD30 / TNFRSF8 Protein (Fc Tag)

Catalog Number: 10777-H02H

General Information

Gene Name Synonym:

CD30; D1S166E; Ki-1

Protein Construction:

A DNA sequence encoding the human TNFRSF8 (NP_001234.2) (Met1-Lys379) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Human

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE.

Endotoxin:

< 1.0 EU per µg protein as determined by the LAL method.

Stability:

Samples are stable for up to twelve months from date of receipt at -70 °C

Predicted N terminal: Phe 19

Molecular Mass:

The recombinant human TNFRSF8 consists of 599 amino acids and predicts a molecular mass of 65.2 kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Storage:


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

CD3, also known as TNFRSF8, is a cell membrane protein of the tumor necrosis factor receptor (TNFR) superfamily. CD3 protein is expressed by activated, but not resting, T and B cells. CD3 can regulate proliferation of lymphocytes and may also play an important role in human immunodeficiency virus replication. As a regulator of apoptosis, CD3 protein induces cell death or proliferation, depending on the cell type, and has been shown to limit the proliferative potential of autoreactive CD8 effector T cells and protect the body against autoimmunity. CD3 protein expression is upregulated in various hematological malignancies, including Reed-Sternberg cells in Hodgkin's disease (HD), anaplastic large cell lymphoma (ALCL) and subsets of Non-Hodgkin's lymphomas (NHLs), and CD3 is also linked to leukocytes in patients with chronic inflammatory diseases, including lupus erythematosus, asthma, rheumatoid arthritis and atopic dermatitis (AD).

References

1. Rossi FM, *et al.* (2001) CD30L up-regulates CD30 and IL-4 expression by T cells. *FEBS Lett.* 508(3): 418-22. 2. Trovato M, *et al.* (2001) Expression of CD30 ligand and CD30 receptor in normal thyroid and benign and malignant thyroid nodules. *Thyroid.* 11(7): 621-8. 3. Ekstrom ES, *et al.* (2001) Presence of CD30(+) and CD30L(+) cells in human placenta and soluble CD30 levels in cord blood are independent of maternal atopy. *Placenta.* 22(4): 372-9.

Manufactured By Sino Biological Inc., FOR RESEARCH USE ONLY. NOT FOR USE IN HUMANS.

For US Customer: Fax: 267-657-0217 ● Tel: 215-583-7898

Global Customer: Fax :+86-10-5862-8288 ● Tel:+86-400-890-9989 ● <http://www.sinobiological.com>