

VISTA/C10orf54 Antibody, Rabbit PAb, Antigen Affinity Purified



Sino Biological
Biological Solution Specialist

Catalog Number: 13482-T26

GENERAL INFORMATION

Immunogen:	Recombinant Human VISTA/C10orf54 Protein (Catalog#13482-H08H)
Preparation	Produced in rabbits immunized with purified, recombinant Human VISTA/C10orf54 (rh VISTA/C10orf54; Catalog#13482-H08H; AAH20568.1; Met1-Ala194). VISTA/C10orf54 specific IgG was purified by Human VISTA/C10orf54 affinity chromatography.
Ig Type:	Rabbit IgG
Specificity:	Human VISTA/C10orf54
Formulation:	0.2 µm filtered solution in PBS
Storage:	This antibody can be stored at 2°C-8°C for one month without detectable loss of activity. Antibody products are stable for twelve months from date of receipt when stored at -20°C to -80°C. Preservative-Free. Avoid repeated freeze-thaw cycles.
Alternative Names:	B7-H5,B7H5,C10orf54,DD1alpha,GI24,PD-1H,PP2135,SISP1,VISTA

APPLICATIONS

Applications:	ELISA,IHC-P,ICC/IF
----------------------	--------------------

RECOMMENDED CONCENTRATION

IHC-P	IHC-P: 1:100-1:500
ICC/IF	ICC/IF: 1:100-1:500
ELISA	ELISA: 1:5000-1:10000 This antibody can be used at 1:5000-1:10000 with the appropriate secondary reagents to detect Human VISTA/C10orf54.


Please Note: Optimal concentrations/dilutions should be determined by the end user.

VISTA/C10orf54 Antibody, Rabbit PAb, Antigen Affinity Purified


Catalog Number: 13482-T26



Sino Biological
Biological Solution Specialist



Immunohistochemical staining of human C10orf54 in human kidney with rabbit polyclonal antibody at 1:200 dilution, formalin-fixed paraffin embedded sections.



Immunofluorescence staining of C10orf54 in U2OS cells. Cells were fixed with 4% PFA, permeabilized with 0.1% Triton X-100 in PBS, blocked with 10% serum, and incubated with rabbit anti-Human C10orf54 polyclonal antibody (dilution ratio 1:200) at 4°C overnight. Then cells were stained with the Alexa Fluor®488-conjugated Goat Anti-rabbit IgG secondary antibody (green). Positive staining was localized to Nucleus and Cytoplasm.