

Rhesus ACE2 / Angiotensin-Converting Enzyme 2 Protein (Fc Tag)

Catalog Number: 90211-C02H



Sino Biological
Biological Solution Specialist

General Information

Gene Name Synonym:

ACE2

Protein Construction:

A DNA sequence encoding the rhesus ACE2 (ACI04564.1) (Met1-Val739) was expressed with the Fc region of human IgG1 at the C-terminus.

Source: Rhesus

Expression Host: HEK293 Cells

QC Testing

Purity: > 95 % as determined by SDS-PAGE

Bio Activity:

Measured by its ability to cleave a fluorogenic peptide substrate, Mca-YVADAPK (Dnp)-OH (Catalog # ES007).

The specific activity is >250 pmol/min/μg.

Measured by its binding ability in a functional ELISA. Immobilized SARS-CoV-2 (2019-nCoV) Spike Protein (S1+S2 ECD, His tag)(40589-V08B1) at 2μg/mL (100μL/well) can bind Rhesus ACE2-Fc, the EC₅₀ of ACE2 is 30-80 ng/mL.

Endotoxin:

< 1.0 EU per μg of the protein as determined by the LAL method

Predicted N terminal: Gln 18

Molecular Mass:

The recombinant Rhesus ACE2 is comprises 963 amino acids and has a calculated molecular mass of 110.7kDa.

Formulation:

Lyophilized from sterile PBS, pH 7.4.

Normally 5 % - 8 % trehalose, mannitol and 0.01% Tween80 are added as protectants before lyophilization. Specific concentrations are included in the hardcopy of COA. Please contact us for any concerns or special requirements.

Usage Guide

Stability & Storage:

Samples are stable for twelve months from date of receipt at -20°C to -80°C.


Store it under sterile conditions at -20°C to -80°C upon receiving. Recommend to aliquot the protein into smaller quantities for optimal storage.

Avoid repeated freeze-thaw cycles.

Reconstitution:

Detailed reconstitution instructions are sent along with the products.

SDS-PAGE:



Protein Description

Angiotensin-converting enzyme 2 (ACE2), a first homolog of ACE, regulates the renin angiotensin system (RAS) by counterbalancing ACE activity. Accumulating evidence in recent years has demonstrated a physiological and pathological role of ACE2 in the cardiovascular, renal and respiratory systems. ACE2 also has an important role in blood pressure control. This enzyme, an homolog of ACE, hydrolyzes angiotensin (Ang) I to produce Ang-(1-9), which is subsequently converted into Ang-(1-7) by a neutral endopeptidase and ACE. ACE2 releases Ang-(1-7) more efficiently than its catalysis of Ang-(1-9) by cleavage of Pro(7)-Phe(8) bound in Ang II. Thus, the major biologically active product of ACE2 is Ang-(1-7), which is considered to be a beneficial peptide of the RAS cascade in the cardiovascular system. A physiological role for ACE2 has been implicated in hypertension, cardiac function, heart function and diabetes, and as a receptor of the severe acute respiratory syndrome coronavirus. In the acute respiratory distress syndrome (ARDS), ACE, AngII, and AT1R promote the disease pathogenesis, whereas ACE2 and the AT2R protect from ARDS. Importantly, ACE2 has been identified as a key SARS-coronavirus receptor and plays a protective role in severe acute respiratory syndrome (SARS) pathogenesis. Furthermore, the recent explosion of research into the ACE2 homolog, collectrin, has revealed a new physiological function of ACE2 as an amino acid transporter, which explains the pathogenic role of gene mutations in Hartnup disorder. This review summarizes and discusses the recently unveiled roles for ACE2 in disease pathogenesis.

References

1. Koitka A, et al. (2008) Angiotensin converting enzyme 2 in the kidney. Clin Exp Pharmacol Physiol. 35(4): 420-5.
2. Raizada MK, et al. (2007) ACE2: a new target for cardiovascular disease therapeutics. J Cardiovasc Pharmacol. 50(2): 112-9.
3. Imai Y, et al. (2007) Angiotensin-converting enzyme 2 (ACE2) in disease pathogenesis. Circ J. 74(3): 405-10.