

## **TECHNICAL DATA SHEET**

# **APC Anti-Human CD10 (SN5c)**

Catalog Number: 20-0108

## PRODUCT INFORMATION

Contents: APC Anti-Human CD10 (SN5c)

Isotype: Mouse IgG1, kappa

Concentration: 5 µL (0.25 µg)/test

Clone: SN5c


Reactivity: Human

**Use By:** 12 months from date of receipt

**Storage Conditions:** 2-8°C protected from light

Formulation: 10 mM NaH2PO4, 150 mM NaCl, 0.09% NaN3,

0.1% gelatin, pH7.2



Human peripheral blood granulocytes were stained with 5 uL (0.25 ug) APC Anti-Human CD10 (20-0108) (solid line) or 0.25 ug APC Mouse IgG1 isotype control (dashed line).

Rev. 20150121

#### **DESCRIPTION**

The SN5c antibody is specific for human CD10, a 100 kDa type II transmembrane protein also known as Common Acute Lymphoblastic Leukemia Antigen (CALLA). In normal cells, it is expressed on early B and T lymphoid precursors, neutrophils and on various epithelia. CD10 is also expressed on several lymphoma cells including Burkitt's, acute lymphoblastic, and follicular germinal center lymphomas. CD10 functions as an endopeptidase and is involved in B cell development, as well as a mediator of neutrophil inflammatory responses.

#### **PREPARATION & STORAGE**

This monoclonal antibody was purified from tissue culture supernatant via affinity chromatography. The purified antibody was conjugated under optimal conditions, with unreacted dye removed from the preparation. It is recommended to store the product undiluted at 4°C, and protected from prolonged exposure to light. Do not freeze.

#### **APPLICATION NOTES**

This antibody preparation has been pre-titrated and quality-tested for flow cytometry using an appropriate cell type. The antibody has been diluted for use at 5  $\mu$ L per test, defined as the amount of antibody that will stain a cell sample in a final volume of approximately 100  $\mu$ L. The number of cells within a sample should be determined empirically, but typically ranges between 1x10e5 to 1x10e8 cells.

### **REFERENCES**

Matsuzaki H, Haruta Y, Fukukawa T, Barcos MP and Seon BK. 1987. Cancer Res. 47(8): 2160-2166.
Shipp MA, Stefano GB, Switzer SN, Griffin JD and Reinherz EL. 1991. Blood. 78(7): 1834-1841.
Pinho S, Lacombe J, Hanoun M, Mizoguchi T, Bruns I, Kunisaki Y and Frenette PS. 2013. J Exp Med. 201(7): 1351-1367. (Flow cytometry) Mukhopadhyay C, Zhao X, Maroni D, Band V and Naramura M. 2013. PLoS One. 8(10): e75907. (Flow cytometry)

Tonbo Biosciences tests all antibodies by flow cytometry. Citations are provided as a resource for additional applications that have not been validated by Tonbo Biosciences. Please choose the appropriate format for each application and consult Materials and Methods sections for additional details about the use of any product in these publications.

For Research Use Only

Not for use in diagnostic or therapeutic procedures. Not for resale. Not for distribution without written consent. Tonbo Biosciences will not be held responsible for patent infringement or other violations that may occur with the use of our products. Tonbo Biosciences, Tonbo Biosciences Logo and all other trademarks are the property of Tonbo Biotechnologies Corporation. © 2013 Tonbo Biosciences.